

BLACK-BRANE SOLUTION FOR C_2 ALGEBRA

M.A. Grebeniuk¹, V.D. Ivashchuk²

*Center for Gravitation and Fundamental Metrology, VNIIMS, 3/1 M.
Ulyanovoy Str., Moscow 117313, Russia and
Institute of Gravitation and Cosmology, People's Friendship University
of Russia, Mikhlukho-Maklaya Str. 6,
Moscow 117198, Russia*

and S.-W. Kim³

*Department of Science Education and Basic Science Research Institute,
Ewha Women's University, Seoul 120-750, Korea*

Abstract

Black p -brane solutions for a wide class of intersection rules and Ricci-flat “internal” spaces are considered. They are defined up to moduli functions H_s obeying non-linear differential equations with certain boundary conditions imposed. A new solution with intersections corresponding to the Lie algebra C_2 is obtained. The functions H_1 and H_2 for this solution are polynomials of degree 3 and 4.

*PACS numbers: 04.20.Jb, 04.50.+h, 04.70.Bw, 02.20.Sv, 02.30.Hq.

¹mag@gravi.phys.msu.su

²ivas@rgs.phys.msu.su

³sungwon@mm.ewha.ac.kr

1 Introduction

The paper is devoted to black-brane solutions with (next to) arbitrary intersections (see Section 2) [1, 2, 3, 4]. These black-brane solutions are governed by moduli functions $H_s = H_s(R)$ obeying a set of second order non-linear differential equations with some boundary relations imposed.

Some general features of the black-brane solutions were investigated earlier in [15]. More general spherically symmetric (and cosmological solutions) were obtained in ref. [5] using the sigma-model approach and the Lagrange representation from refs. [6, 13].

In [1] the following conjecture was suggested (see Section 3): the moduli functions H_s are polynomials when intersection rules correspond to semisimple Lie algebras. This conjecture was confirmed by special black-brane "block-orthogonal" solutions considered earlier in [14, 16, 18] (subsect. 3.1). It was verified also for A_m and C_m series of Lie algebras in [2, 3]. An analogue of this conjecture for extremal black holes was considered earlier in [11].

An example of black brane solution corresponding to Lie algebra $A_2 = sl(3)$ was considered earlier in refs. [1, 2], where some dyonic configurations (e.g. in 11-dimensional supergravity) were considered. The A_2 -solution is governed by two polynomials H_1 and H_2 of degree 2. The coefficients of polynomials and charges are functions of some parameters P_1 and P_2 and have a rather simple form (subsect. 3.1). It seems that A_2 -solution is the only one that may be obtained "by hands": the search of solutions for other Lie algebras needs computer calculations.

Here we present a first non-trivial result of such calculations, i.e. a new solution corresponding to the Lie algebra C_2 governed by polynomials of degree 3 and 4 (subsect. 3.3). The coefficients of polynomials exhibit a non-trivial dependence upon parameters P_1 and P_2 , namely: a non-trivial term Δ , that is a square root of a polynomial of degree 4, appears.

2 p -brane black hole solutions

Consider a model governed by the action

$$S = \int d^D x \sqrt{|g|} \left\{ R[g] - h_{\alpha\beta} g^{MN} \partial_M \varphi^\alpha \partial_N \varphi^\beta - \sum_{a \in \Delta} \frac{\theta_a}{n_a!} \exp[2\lambda_a(\varphi)] (F^a)^2 \right\} \quad (1)$$

where $g = g_{MN}(x)dx^M \otimes dx^N$ is a metric, $\varphi = (\varphi^\alpha) \in \mathbb{R}^l$ is a vector of scalar fields, $(h_{\alpha\beta})$ is a constant symmetric non-degenerate $l \times l$ matrix ($l \in \mathbb{N}$), $\theta_a = \pm 1$,

$$F^a = dA^a = \frac{1}{n_a!} F_{M_1 \dots M_{n_a}}^a dz^{M_1} \wedge \dots \wedge dz^{M_{n_a}} \quad (2)$$

is a n_a -form ($n_a \geq 1$), λ_a is a 1-form on $\mathbb{R}^l$: $\lambda_a(\varphi) = \lambda_{\alpha a} \varphi^\alpha$, $a \in \Delta$, $\alpha = 1, \dots, l$. In (1) we denote $|g| = |\det(g_{MN})|$, $(F^a)_g^2 = F_{M_1 \dots M_{n_a}}^a F_{N_1 \dots N_{n_a}}^a g^{M_1 N_1} \dots g^{M_{n_a} N_{n_a}}$, $a \in \Delta$. Here Δ is some finite set. In the models with one time all $\theta_a = 1$ when the signature of the metric is $(-1, +1, \dots, +1)$.

Let us consider (black-brane) solutions to field equations corresponding to the action (1) from [1, 2, 3]. These solutions are defined on the manifold

$$M = (R_0, +\infty) \times (M_1 = S^{d_1}) \times (M_2 = \mathbb{R}) \times \dots \times M_n, \quad (3)$$

and have the following form

$$g = \left(\prod_{s \in S} H_s^{2h_s d(I_s)/(D-2)} \right) \left\{ f^{-1} dR \otimes dR + R^2 d\Omega_{d_1}^2 \right. \quad (4)$$

$$\left. - \left(\prod_{s \in S} H_s^{-2h_s} \right) f dt \otimes dt + \sum_{i=3}^n \left(\prod_{s \in S} H_s^{-2h_s \delta_{iI_s}} \right) g^i \right\},$$

$$\exp(\varphi^\alpha) = \prod_{s \in S} H_s^{h_s \chi_s \lambda_{a_s}^\alpha}, \quad (5)$$

$$F^a = \sum_{s \in S} \delta_{a_s}^a \mathcal{F}^s, \quad (6)$$

where $f = 1 - 2\mu/R^{\bar{d}}$,

$$\mathcal{F}^s = -\frac{Q_s}{R^{d_1}} \left(\prod_{s' \in S} H_{s'}^{-A_{ss'}} \right) dR \wedge \tau(I_s), \quad s \in S_e, \quad (7)$$

$$\mathcal{F}^s = Q_s \tau(\bar{I}_s), \quad s \in S_m. \quad (8)$$

Here $Q_s \neq 0$ ($s \in S$) are charges, $R_0 > 0$, $R_0^{\bar{d}} = 2\mu > 0$, $\bar{d} = d_1 - 1$. In (4) $g^i = g_{m_i n_i}^i(y_i) dy_i^{m_i} \otimes dy_i^{n_i}$ is a Ricci-flat metric on M_i , $i = 3, \dots, n$ and $\delta_{iI} = \sum_{j \in I} \delta_{ij}$ is the indicator of i belonging to I : $\delta_{iI} = 1$ for $i \in I$ and

$\delta_{iI} = 0$ otherwise. Here $g^2 = -dt \otimes dt$, and $g^1 = d\Omega_{d_1}$ be a canonical metric on unit sphere $M_1 = S^{d_1}$,

The p -brane set S is by definition

$$S = S_e \cup S_m, \quad S_v = \cup_{a \in \Delta} \{a\} \times \{v\} \times \Omega_{a,v}, \quad (9)$$

$v = e, m$ and $\Omega_{a,e}, \Omega_{a,m} \subset \Omega$, where $\Omega = \Omega(n)$ is the set of all non-empty subsets of $\{2, \dots, n\}$, i.e. all p -branes do not “live” in M_1 .

Any p -brane index $s \in S$ has the form $s = (a_s, v_s, I_s)$, where $a_s \in \Delta$, $v_s = e, m$ and $I_s \in \Omega_{a_s, v_s}$. The sets S_e and S_m define electric and magnetic p -branes, correspondingly. In (5) $\chi_s = +1, -1$ for $s \in S_e, S_m$, respectively. All p -branes contain the time manifold $M_2 = \mathbb{R}$, i.e.

$$2 \in I_s, \quad \forall s \in S. \quad (10)$$

All manifolds M_i , $i > 2$, are oriented and connected and

$$\tau_i \equiv \sqrt{|g^i(y_i)|} dy_i^1 \wedge \dots \wedge dy_i^{d_i}, \quad (11)$$

are volume d_i -forms, where $d_i = \dim M_i$, $i = 1, \dots, n$, with $d_1 > 1$ and $d_2 = 1$. For any $I = \{i_1, \dots, i_k\} \in \Omega$, $i_1 < \dots < i_k$, we denote

$$\tau(I) \equiv \tau_{i_1} \wedge \dots \wedge \tau_{i_k}, \quad d(I) = \sum_{i \in I} d_i. \quad (12)$$

The forms $\mathcal{F}^s$ correspond to electric and magnetic p -branes for $s \in S_e, S_m$, respectively. In (8) we use the notation $\bar{I} = \{1, \dots, n\} \setminus I$.

The parameters h_s appearing in the solution satisfy the relations: $h_s = (B_{ss})^{-1}$, where

$$B_{ss'} = d(I_s \cap I_{s'}) + \frac{d(I_s)d(I_{s'})}{2 - D} + \chi_s \chi_{s'} \lambda_{\alpha a_s} \lambda_{\beta a_{s'}} h^{\alpha\beta}, \quad (13)$$

$s, s' \in S$, with $(h^{\alpha\beta}) = (h_{\alpha\beta})^{-1}$ and $D = 1 + \sum_{i=1}^n d_i$. Here we assume that (i) $B_{ss} \neq 0$, $s \in S$, and (ii) $\det(B_{ss'}) \neq 0$, i.e. the matrix $(B_{ss'})$ is a non-degenerate one.

Let us consider the matrix

$$(A_{ss'}) = (2B_{ss'}/B_{s's'}). \quad (14)$$

Here some ordering in S is assumed.

The functions $H_s = H_s(z) > 0$, $z = 2\mu/R^{\bar{d}} \in (0, 1)$, obey the equations

$$\frac{d}{dz} \left(\frac{(1-z)}{H_s} \frac{dH_s}{dz} \right) = B_s \prod_{s' \in S} H_{s'}^{-A_{ss'}}, \quad (15)$$

equipped with the boundary conditions

$$H_s(1-0) = H_{s0} \in (0, +\infty), \quad (16)$$

$$H_s(+0) = 1, \quad (17)$$

$s \in S$. Here $B_s = B_{ss}\varepsilon_s Q_s^2 / (2\bar{d}\mu)^2$ and

$$\varepsilon_s = (-\varepsilon[g])^{(1-\chi_s)/2} \varepsilon(I_s) \theta_{a_s}, \quad (18)$$

$s \in S$, $\varepsilon[g] \equiv \text{sign det}(g_{MN})$. More explicitly eq. (18) reads: $\varepsilon_s = \varepsilon(I_s) \theta_{a_s}$ for $v_s = e$ and $\varepsilon_s = -\varepsilon[g] \varepsilon(I_s) \theta_{a_s}$, for $v_s = m$.

Equations (16) are equivalent to Toda-type equations. The first boundary condition guarantees the existence of a regular horizon at $R^{\bar{d}} = 2\mu$. The second condition (17) ensures an asymptotical flatness (for $R \rightarrow +\infty$) of the $(2+d_1)$ -dimensional section of the metric.

Due to eqs. (7) and (8), the dimension of p -brane worldvolume $d(I_s)$ is defined by relations: $d(I_s) = n_{a_s} - 1$, $d(I_s) = D - n_{a_s} - 1$, for $s \in S_e, S_m$, respectively. For a p -brane we use a standard notation: $p = p_s = d(I_s) - 1$.

The solutions are valid if the following restriction on the sets $\Omega_{a,v}$ is imposed. This restriction guarantees the block-diagonal structure of the stress-energy tensor. We denote $w_1 \equiv \{i | i \in \{2, \dots, n\}, d_i = 1\}$ and $n_1 = |w_1|$ (i.e. n_1 is the number of 1-dimensional spaces among M_i , $i = 2, \dots, n$). It is clear, that $2 \in w_1$.

Restriction. *Let 1a) $n_1 \leq 1$ or 1b) $n_1 \geq 2$ and for any $a \in \Delta$, $v \in \{e, m\}$, $i, j \in w_1$, $i \neq j$, there are no $I, J \in \Omega_{a,v}$ such that $i \in I$, $j \in J$ and $I \setminus \{i\} = J \setminus \{j\}$.*

This restriction is satisfied in the non-composite case: $|\Omega_{a,e}| + |\Omega_{a,m}| = 1$, (i.e when there are no two p -branes with the same color index a , $a \in \Delta$.) The restriction forbids certain intersections of two p -branes with the same color index for $n_1 \geq 2$.

The solution under consideration describes a set of charged (by forms) overlapping black p -branes "living" on submanifolds of $M_2 \times \dots \times M_n$.

3 Examples of solutions

3.1 “Block-orthogonal” solutions

The simplest polynomial solutions occur in orthogonal case [8, 9, 10, 12, 13], when

$$B_{ss'} = 0, \quad (19)$$

for $s \neq s'$, $s, s' \in S$. In this case $(A_{ss'}) = \text{diag}(2, \dots, 2)$ is a Cartan matrix for semisimple Lie algebra $A_1 \oplus \dots \oplus A_1$ and

$$H_s(z) = 1 + P_s z, \quad (20)$$

with $P_s \neq 0$, satisfying

$$P_s(P_s + 1) = -B_s, \quad (21)$$

$s \in S$. For positive parameters $P_s > 0$ we get negative $B_s < 0$.

In [14, 16, 18] this solution was generalized to the “block-orthogonal” case:

$$S = S_1 \cup \dots \cup S_k, \quad S_i \cap S_j = \emptyset, \quad i \neq j, \quad (22)$$

$S_i \neq \emptyset$, i.e. the set S is a union of k non-intersecting (non-empty) subsets $S_1, \dots, S_k$, and relation (19) is satisfied for all $s \in S_i$, $s' \in S_j$, $i \neq j$; $i, j = 1, \dots, k$. In this case eq. (20) is modified as follows

$$H_s(z) = (1 + P_s z)^{b^s}, \quad (23)$$

where

$$b^s = 2 \sum_{s' \in S} A^{ss'}, \quad (24)$$

$(A^{ss'}) = (A_{ss'})^{-1}$ and parameters P_s are coinciding inside blocks, i.e. $P_s = P_{s'}$ for $s, s' \in S_i$, $i = 1, \dots, k$. Parameters $P_s \neq 0$ satisfy the relations (21) and parameters B_s are also coinciding inside blocks, i.e. $B_s = B_{s'}$ for $s, s' \in S_i$, $i = 1, \dots, k$.

Let $(A_{ss'})$ be a Cartan matrix for a finite-dimensional semisimple Lie algebra $\mathcal{G}$. In this case all powers in (24) are natural numbers coinciding with the components of twice the dual Weyl vector in the basis of simple roots [19], and hence, all functions H_s are polynomials, $s \in S$.

Conjecture [1]. *Let $(A_{ss'})$ be a Cartan matrix for a semisimple finite-dimensional Lie algebra $\mathcal{G}$. Then the solution to eqs. (15)-(17) (if exists) is a polynomial*

$$H_s(z) = 1 + \sum_{k=1}^{n_s} P_s^{(k)} z^k, \quad (25)$$

where $P_s^{(k)}$ are constants, $k = 1, \dots, n_s$, the integers $n_s = b^s$ are defined in (24) and $P_s^{(n_s)} \neq 0$, $s \in S$.

This conjecture was verified for A_n and C_n series of Lie algebras [2, 3]. In the extremal case $\mu = +0$ an analogue of this conjecture was suggested previously in [11].

3.2 Solution for A_2 algebra

Here we present the polynomial solution from [1, 2] corresponding to the Lie algebra $A_2 = sl(3)$ with the Cartan matrix

$$(A_{ss'}) = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} \quad (26)$$

The moduli polynomials read in this case as follows

$$H_s = 1 + P_s z + P_s^{(2)} z^2, \quad (27)$$

where $P_s = P_s^{(1)}$ and $P_s^{(2)} \neq 0$ are constants and

$$P_s^{(2)} = \frac{P_s P_{s+1} (P_s + 1)}{2(P_1 + P_2 + 2)}, \quad (28)$$

$$B_s = -\frac{P_s (P_s + 1) (P_s + 2)}{P_1 + P_2 + 2}, \quad (29)$$

$s = 1, 2$. Here $P_1 + P_2 + 2 \neq 0$.

In the A_2 -case the solution is described by relations (4)-(8) with $S = \{s_1, s_2\}$ and intersection rules following from (13), (14) and (26)

$$d(I_{s_1} \cap I_{s_2}) = \frac{d(I_{s_1})d(I_{s_2})}{D-2} - \chi_{s_1} \chi_{s_2} \lambda_{a_{s_1}} \cdot \lambda_{a_{s_2}} - \frac{1}{2}K, \quad (30)$$

$$d(I_{s_i}) - \frac{(d(I_{s_i}))^2}{D-2} + \lambda_{a_{s_i}} \cdot \lambda_{a_{s_i}} = K, \quad (31)$$

where $K \neq 0$ and functions $H_{s_i} = H_i$ are defined by relations (27)-(29) with $z = 2\mu R^{-\vec{d}}$, $i = 1, 2$. Here and in what follows $\lambda \cdot \lambda' = \lambda_\alpha \lambda'_\beta h^{\alpha\beta}$.

3.3 Solutions for C_2 algebra

Now we present the solution related to the Lie algebra $C_2 = so(5)$ with the Cartan matrix

$$(A_{ss'}) = \begin{pmatrix} 2 & -1 \\ -2 & 2 \end{pmatrix}. \quad (32)$$

According to **Conjecture** we seek the solution to eqs. (15)-(17) in the following form

$$H_1(z) = 1 + P_1 z + P_1^{(2)} z^2 + P_1^{(3)} z^3, \quad (33)$$

$$H_2(z) = 1 + P_2 z + P_2^{(2)} z^2 + P_2^{(3)} z^3 + P_2^{(4)} z^4, \quad (34)$$

where $P_s = P_s^{(1)}$ and $P_s^{(k)}$ are constants, $s = 1, 2$.

Here we outline the result. For B_s -parameters we get the following relations

$$2B_1 = -\Delta + (2P_1 + 3)(2 + P_2), \quad (35)$$

$$B_2 = \Delta - 2 - 2P_1(P_1 + 3) - (2 + P_2)^2, \quad (36)$$

and for parameters $P_s^{(k)}$ we obtain

$$4P_1^{(2)} = 6 + 3P_2 - \Delta + 2P_1(3 + P_1 + P_2), \quad (37)$$

$$12P_1^{(3)} = -\Delta(2 + P_1 + P_2) + 12 + 18P_1 + 2P_1^3 + 3P_2(4 + P_2) + 2P_1^2(5 + P_2) + P_1P_2(11 + 2P_2), \quad (38)$$

$$2P_2^{(2)} = -6 - 2P_1(3 + P_1) - 3P_2 + \Delta, \quad (39)$$

$$6P_2^{(3)} = \Delta(2 + 2P_1 + P_2) - 12 - 24P_1 - 4P_1^3 - 3P_2(4 + P_2) - 2P_1P_2(7 + P_2) - 2P_1^2(8 + P_2), \quad (40)$$

$$24P_2^{(4)} = \Delta[2P_1^2 + (3 + P_2)(2 + 2P_1 + P_2)] - 4P_1^4 - 3(2 + P_2)^2(3 + P_2) - 2P_1(3 + P_2)^2(4 + P_2) - 4P_1^3(6 + P_2) - P_1^2(60 + 30P_2 + 4P_2^2), \quad (41)$$

where

$$\Delta = \sqrt{4(3 + P_1(3 + P_1))^2 + (3 + 2P_1)^2 P_2(4 + P_2)}. \quad (42)$$

It may be verified that $B_1 < 0$ and $B_2 < 0$ for $P_1 > 0$, $P_2 > 0$.

The C_2 black-brane solution is described by relations (4)-(8), with $S = \{s_1, s_2\}$, and intersection rules following from (13), (14) and (32)

$$d(I_{s_1} \cap I_{s_2}) = \frac{d(I_{s_1})d(I_{s_2})}{D-2} - \chi_{s_1}\chi_{s_2}\lambda_{a_{s_1}} \cdot \lambda_{a_{s_2}} - K, \quad (43)$$

$$d(I_{s_1}) - \frac{(d(I_{s_1}))^2}{D-2} + \lambda_{a_{s_1}} \cdot \lambda_{a_{s_1}} = K, \quad (44)$$

$$d(I_{s_2}) - \frac{(d(I_{s_2}))^2}{D-2} + \lambda_{a_{s_2}} \cdot \lambda_{a_{s_2}} = 2K, \quad (45)$$

where $K \neq 0$ and functions $H_{s_i} = H_i$ are defined by relations (33) and (34) with $z = 2\mu R^{-\bar{d}}$, $i = 1, 2$.

A simple test for verification of the relations for H_s -functions is to put $P_1 = 3P$ and $P_2 = 4P$ with $P > 0$. In this case we get a special “block-orthogonal” solution

$$H_1(z) = (1 + Pz)^3, \quad H_2(z) = (1 + Pz)^4, \quad (46)$$

in agreement with the relations (23) and (24).

4 Conclusions

In this paper we presented a new black-brane solution with two branes and intersection rule corresponding to the Lie algebra C_2 . The solutions is governed by polynomials of degree 3 and 4 and the coefficients of polynomials exhibit a non-trivial dependence upon the parameters P_1 and P_2 , due to appearance of the non-trivial term Δ (see (42)). The C_2 -solution differs drastically from the A_2 -one, that has a rather simple analytical structure. This means that the polynomial solutions for other A_n -algebras ($n > 2$) should be also non-trivial. Indeed, the C_2 -solution may be extended to a special A_3 -one, if we impose a restriction on the moduli polynomials H_1, H_2, H_3 of the following form: $H_1 = H_3$. (Thus, here we obtained by product a special A_3 black-brane solution.) It should be noted that A, D, E (or simply laced) Lie algebras are of much interest, since they appear for p -brane intersection rules in supergravitational models [13]. Another topics of interest are related to black-brane thermodynamics (e.g. relations for the entropy, the Hawking temperature etc) and analysis of post-Newtonian effects. On this way one

may expect to clarify the appearance of the function Δ in the solution. But the main (mathematical) problem here is to find the polynomial solutions for all Lie algebras. This problem seems to be a very difficult one and may be of interest for mathematicians dealing with polynomials (e.g. appearing in non-linear ordinary and partial differential equations), Lie algebras, number theory etc.

Acknowledgments

This work was supported in part by the Russian Ministry for Science and Technology, Russian Foundation for Basic Research, project SEE, by grant No. R01-2000-00015 from the Korea Science and Engineering Foundation and the MOST through National Research Program (00-B-WB-06-A-04) for Women's University. V.D.I. thanks the colleagues from Ewha Women's University for kind hospitality. The results of the work were reported at the workshop "International Camp of Gravitation" (Sept. 8-9, 2001, Institute of Wind and Water, Gapyong, Korea) and at Fifth International conference on Gravitation and Astrophysics of Asia-Pacific countries.

References

- [1] V.D. Ivashchuk and V.N. Melnikov, P-brane black holes for general intersections, *Gravitation & Cosmology* **5**, 313-318 (1999); gr-qc/0002085.
- [2] V.D. Ivashchuk and V.N. Melnikov, Black hole p-brane solutions for general intersection rules. *Gravitation & Cosmology* **6**, 27-40 (2000); hep-th/9910041.
- [3] V.D. Ivashchuk and V.N. Melnikov, Toda p-brane black holes and polynomials related to Lie algebras. *Class. Quantum Grav.* **17**, 2073-2092 (2000); math-ph/0002048.
- [4] V.D. Ivashchuk and V.N. Melnikov, Exact solutions in multidimensional gravity with antisymmetric forms, topical review, *Class. Quantum Grav.* **18**, R1-R66 (2001); hep-th/0110274.
- [5] V.D.Ivashchuk and S.-W. Kim, Solutions with intersecting p-branes related to Toda chains, *J. Math. Phys.* **41**, 444-460 (2000); hep-th/9907019.

- [6] V.D. Ivashchuk and V.N. Melnikov, Sigma-model for the generalized composite p-branes, *Class. Quantum Grav.* **14**, 3001-3029 (1997); *Corrigenda* **15**, 3941 (1998); hep-th/9705036.
- [7] V.D.Ivashchuk and V.N.Melnikov, Exact solutions in sigma-model with p-branes, *J. Korean Phys. Soc.* **35**, S638-S648 (1999).
- [8] M. Cvetič and A. Tseytlin, Non-extreme black holes from non-extreme intersecting M -branes, *Nucl. Phys. B* **478**, 181-198 (1996); hep-th/9606033.
- [9] I.Ya. Aref'eva, M.G. Ivanov and I.V. Volovich, Non-extremal intersecting p-branes in various dimensions, *Phys. Lett. B* **406**, 44-48 (1997); hep-th/9702079.
- [10] N. Ohta, Intersection rules for non-extreme p-branes, *Phys. Lett. B* **403**, 218-224 (1997); hep-th/9702164.
- [11] H. Lü, J. Maharana, S. Mukherji and C.N. Pope, Cosmological solutions, p-branes and the Wheeler-DeWitt equation, *Phys. Rev. D* **57**, 2219-2229 (1998); hep-th/9707182.
- [12] K.A. Bronnikov, V.D. Ivashchuk and V.N. Melnikov, The Reissner-Nordström problem for intersecting electric and magnetic p-branes, *Gravitation & Cosmology* **3**, 203-212 (1997); gr-qc/9710054.
- [13] V.D. Ivashchuk and V.N. Melnikov, Multidimensional classical and quantum cosmology with intersecting p -branes, *J. Math. Phys.* **39**, 2866-2889 (1998); hep-th/9708157.
- [14] K.A. Bronnikov, Block-orthogonal brane systems, black holes and wormholes, *Gravitation & Cosmology* **4**, 49-56 (1998); hep-th/9710207.
- [15] K.A. Bronnikov, Gravitating brane systems: some general theorems, *J. Math. Phys.* **40**, 2, 924-938 (1999); gr-qc/9806102.
- [16] V.D. Ivashchuk and V.N. Melnikov. Multidimensional cosmological and spherically symmetric solutions with intersecting p-branes. In Lecture

Notes in Physics, Vol. 537, *Mathematical and Quantum Aspects of Relativity and Cosmology*, Proceedings of the Second Samos Meeting on Cosmology, Geometry and Relativity held at Pythagoreon, Samos, Greece, 1998, eds. S. Cotsakis, G.W. Gibbons (Springer, Berlin, 2000); gr-qc/9901001.

- [17] V.D.Ivashchuk and V.N.Melnikov, Cosmological and spherically symmetric solutions with intersecting p-branes. *J. Math. Phys.* **40**, 6558-6576 (1999).
- [18] S. Cotsakis, V.D. Ivashchuk and V.N. Melnikov, P-branes black holes and post-Newtonian approximation, *Gravitation & Cosmology* **5**, 52-57 (1999); gr-qc/9902148.
- [19] J. Fuchs and C. Schweigert, *Symmetries, Lie algebras and representations. A graduate course for physicists* (Cambridge University Press, Cambridge, 1997).